

# FLOWERING SEASONS OF AUSTRALIAN PLANTS.

BY E. HAVILAND, F.L.S.

## NO. I.—LIST OF PLANTS INDIGENOUS IN THE NEIGHBOURHOOD OF SYDNEY, FLOWERING DURING JULY.

As it is impossible for the author to visit more than a few localities during each month, these lists must necessarily, for the present, be very imperfect. Supplementary lists will, however, be added as each month recurs, until, it is hoped, the flowering seasons of at least all the Cumberland plants have been obtained.

### Dilleniaceæ—

*Hibbertia stricta*  
 „ *acicularis*  
 „ *linearis*  
 „ *Billardieri*.

### Pittosporæ—

*Billardiera scandens*.

### Violaceæ—

*Viola hederacea*  
*Hybanthus Vernonii*

### Rutaceæ—

*Eriostemon lanceolatus*  
 „ *Crowei*  
 „ *buxifolius*  
*Phebalium squamulosum*  
*Philotheca australis*  
*Zieria Smithii*  
 „ *laevigata*  
 „ *pilosa*

### Rutaceæ—

*Boronia ledifolia*  
 „ *anemonifolia*  
 „ *polygalifolia*  
*Correa alba*  
 „ *speciosa*.

### Droseraceæ—

*Drosera peltata*

### Geraniaceæ—

*Pelargonium australe*  
*Oxalis corniculata*.

### Sterculiaceæ—

*Lasiopetalum ferrugineum*.

### Euphorbiaceæ—

*Ricinocarpus pinifolius*  
*Poranthera corymbosa*.

### Leguminosæ—

*Hardenbergia monophylla*  
*Kennedyia rubicunda*

## Leguminosæ—

*Kennedyia prostrata**Hovea linearis*,, *longifolia**Dillwynia ericifolia*,, *floribunda**Aotus villosa**Platylobium formosum**Glycine clandestina**Bossicea heterophylla*,, *rhombofolia*,, *scolopendria**Acacia myrtifolia*,, *suaveolens*,, *longifolia.*,, *oxycedrus*,, *discolor*,, *juniperina*,, ,, var. *Brownei**Gompholobium pinnatum*,, *grandiflorum*,, *glabratum**Pultencea elliptica*,, *stipularis*,, *daphnoides*,, *incurvata*

## Saxifrageæ—

*Bauera rubioides.*

## Halorageæ—

*Haloragis salsoloides.*

## Myrtaceæ—

*Darwinia fascicularis*,, *taxifolia**Callistemon lanceolatus**Calythrix tetragona**Leptospermum flavescens.*

## Rhamnaceæ—

*Cryptandra amara.*

## Umbellifereæ—

*Actinotus minor**Siebera linearifolia**Xanthosia tridentata.*

## Proteaceæ—

*Persoonia lanceolata**Grevillea linearis*,, *sericea*,, *punicea**Conospermum ericifolium*,, *longifolium*,, *taxifolium**Symphyonema paludosum**Banksia ericifolia.*

## Thymeleæ—

*Pimelea linifolia.*

## Rubiaceæ—

*Opercularia aspera**Pomax umbellata.*

## Compositæ—

*Cotula coronopifolia**Aster ramulosus**Brachycome linearifolia**Cassinia denticulata**Senecio lautus.*

## Goodeniaceæ—

*Goodenia hederacea*,, *heterophylla**Dampiera stricta**Velleia lyrata.*

## Loganiaceæ—

*Logania floribunda**Mitrasacme polymorpha.*

## Solanaceæ—

*Solanum nigrum.*

## Labiataæ—

*Hemigenia purpurea**Westringia rosmarinifolia.*

## Verbenaceæ—

*Chloanthes Stœchadis.*

## Epacrideæ—

*Epacris longiflora*,, *obtusifolia*,, *microphylla**Styphelia tubiflora*,, *viridis*,, *longifolia**Leucopogon ericoides*,, *juniperinus*

## Epacrideæ—

*Leucopogon microphyllus**Woolisia pungens**Monotoca elliptica**Sprengelia incarnata.*

## Irideæ—

*Patersonia longifolia*,, *sericea*

## Orchideæ—

*Diuris aurea*,, *maculata**Caladenia alba**Glossodia major**Pterostylis nutans*,, *grandiflora**Prasophyllum fimbriatum.*

All the plants enumerated above, excepting *Acacia oxycedrus*, may be found on the coast line in the immediate neighbourhood of Sydney.